

University of Groningen

Targeted apoptosis induction in hemato-oncology

Cate, Bram ten

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2010

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Cate, B. T. (2010). *Targeted apoptosis induction in hemato-oncology*. s.n.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Targeted elimination of leukemia stem cells; a new therapeutic approach in hemato-oncology

B. ten Cate¹, M. de Bruyn¹, Y. Wei^{1,2}, E. Bremer¹
and W. Helfrich^{1,2}

¹Department of Surgery, Surgical Research Laboratories, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands ²Department of General Surgery, the First Clinical Hospital of Harbin Medical University, Harbin, China

7

Current Drug Targets 2010 January;11(1):95-110

ABSTRACT

Despite recent advances, treatment of leukemia is often not curative. New insights indicate that this may be attributable to a small population of therapy-resistant malignant cells with self-renewal capacity and the ability to generate large numbers of more differentiated leukemia cells. These leukemia-initiating cells are commonly referred to as Leukemia Stem Cells (LSCs). LSCs are regarded as the root of leukemia origin and leukemia recurrence after seemingly successful therapy. Not surprisingly therefore, contemporary leukemia research has focused on ways to specifically eliminate LSCs, leading to the identification of several promising anti-LSC strategies. Firstly, LSCs may be eliminated by antibody- or ligand-based cell surface delivery of therapeutics such as naked antibodies, immunotoxins, and immunocytokines. This approach exploits LSC-associated surface antigens, such as CD33, CD44, CD96, CD123 and CLL-1 for LSC-selective therapy and aims to spare normal hematopoietic stem cells. A second strategy aims to disrupt the interactions between LSCs and their highly specialized niche. These interactions appear to be pivotal for maintenance of the stem cell-like characteristics of LSCs. A third strategy centers on the selective modulation of aberrantly activated signaling pathways central to LSC biology. A fourth strategy, dubbed 'epigenetic reprogramming', aims to selectively reverse epigenetic alterations that are implicated in ontogeny and maintenance of LSCs. In this review, we will discuss the rationale for these LSCs-targeted strategies and highlight recent advances that may ultimately help pave the way towards selective LSCs-elimination.

INTRODUCTION

To a varying degree most leukemia types initially respond well to therapy with partial or even complete remissions. Unfortunately, after a period of minimal residual disease (MRD) many patients succumb to refractory relapses of the disease. Recent insight indicates that the development of these relapses may be due to the selective continued survival of a small, but distinct, population of therapy-resistant tumor-initiating cells, commonly referred to as Leukemia Stem Cells (LSCs). LSCs are thought to originate either from normal hematopoietic stem cells (HSCs) or from more differentiated progenitor cells that have acquired malignant features. In the latter case, the progenitor cells have de-differentiated and re-acquired stem cell-like characteristics via as yet undefined pathways [1;2]. Both LSCs and HSCs possess self-renewal capacity but are relative quiescent compared to more mature progenitors cells (see Fig. 1). However, LSCs typically have a much stronger capacity for cellular expansion than normal HSCs, probably due to an increase in symmetric self-renewal activity of LSCs [3].

LSCs were first identified in Acute Myeloid Leukemia (AML) by John Dick and colleagues [3;4] and have by now been found in all major forms of leukemia [3-6]. LSCs are experimentally defined as cells capable of serial engraftment in immunocompromised mice (see Fig. 1). Importantly, mice engrafted with these LSCs develop leukemia which resembles the patient's original disease [7]. During this last decade substantial progress in the field of LSC research has led to the identification of many of the hallmark

characteristics of LSCs. Therapy-resistance is typical for LSCs which is thought to arise from (1) their relative quiescent cellular activity [8;9], (2) their array of self-protecting mechanisms [10], and (3) the highly protective microenvironment in which the LSCs resides [11].

Since curative treatment of leukemia will most likely require the elimination of LSCs, new strategies need to be designed that overcome the above-described features. Importantly, in this design care should be taken to spare normal HSCs. In this respect, strategies that target a distinct Achilles' heel of the LSC biology appear to be particularly promising. Here, we review the rationale of different LSCs-specific strategies and highlight recent advances that may ultimately lead to selective LSCs-elimination.




Figure 1. Hypotheses for the ontogeny of leukemia stem cells. LSCs can arise by somatic mutations in normal hematopoietic stem cells (HSC) or more mature progenitor cells can become malignant, de-differentiate and re-acquire stem cell-like characteristics. Currently, LSCs (typically CD34⁺CD38⁻) are defined by the ability to engraft in immunocompromised mice and initiate leukemia resembling the original patient's disease.

LSC CELL SURFACE TARGETING APPROACHES

Cell surface antigens expressed on LSCs may be exploited for antibody- or ligand-based therapeutic approaches. The ideal target antigen is exclusively and abundantly expressed on LSCs, thereby maximizing the therapeutic effect and minimizing off-target effects. Unfortunately, such an antigen has not been uncovered. However, a series of surrogate cell surface markers have been identified which have favorable LSCs-restricted expression profiles. These surrogate markers can be targeted with naked antibodies, which typically eliminate targeted cells via Antibody Dependent Cellular Cytotoxicity (ADCC) or Complement Dependent Cytotoxicity (CDC). However, some antibodies can directly interfere with the signaling activity by binding to receptor molecules and sensitize targeted cells to apoptosis induction [12] or directly induce apoptosis as reported for the anti-CD20 monoclonal antibody (mAb) Rituximab [13]. The tumoricidal potential of antibodies can be further enhanced by conjugating them to cytotoxic drugs or radionuclides. Alternatively, cell surface molecules may also be targeted using recombinant forms of their corresponding natural ligand. Ligand-based approaches have been used to deliver cytotoxic drugs to the target cell and to trigger apoptosis-inducing receptors. However, an obvious potential limitation to antibody-based and ligand-based therapy is the selective loss of the target antigen during therapy or the presence of target antigen-negative leukemia cells at the start of therapy. In this respect, fluctuations in the expression of cell surface markers on HSCs have been reported [14]. Therefore, the rational design of combinatorial antibody-based or ligand-based strategies that target two or more different LSC antigens simultaneously may prove the best way to prevent escape from therapy. Below, a selection of the most prominent cell surface molecules on LSCs and approaches for targeting these molecules are briefly reviewed (summarized in Fig. 2).

LSC Target antigens

CD33 belongs to immunoglobulin (Ig) superfamily and is a member of the sialoadhesin family of cellular interaction molecules [15;16]. The intracellular domain of CD33 contains an immunoreceptor tyrosine-based inhibitory motif (ITIM) domain which may function as a negative regulator of cellular differentiation [17]. CD33 expression is largely restricted to the myeloid lineage, however some expression is detected on certain peripheral blood lymphocytes. In leukemia, CD33 is expressed on blasts in the majority of AML patients. Importantly, it has been noted that CD33 is also expressed on AML-LSCs, but not on normal HSCs [18;19].

CD123 (IL-3R α) is the alpha subunit that hetero-dimerizes with the IL3 β chain to form the interleukin-3 receptor. CD123 is constitutively expressed on hematopoietic progenitors cells (HPCs), on which it is involved in proliferation and differentiation. Furthermore, CD123 is expressed on LSCs in AML [20], Chronic Myeloid Leukemia (CML), Myelodysplastic Syndrome (MDS) and Systemic Mastocytosis [18], but not on HSCs [21]. This expression profile indicates that CD123 might be a suitable target for the selective elimination of LSCs in various hematological malignancies.

C-type Lectin-Like Molecule-1 (CLL-1) is a heavily N-glycosylated transmembrane protein with an intracellular ITIM domain [22]. CLL-1 is exclusively expressed on cells of the hematopoietic lineage, including dendritic cells, granulocytes and monocytes [23]. In leukemia, CLL-1 expression is detected on 92% of all AML cases [23] on which a high expression at diagnosis correlates with an early relapse [24]. Importantly, CLL-1 is also expressed on AML-LSCs, but not on HSCs [24;25]. In line with these findings, it was demonstrated that sorted CD34⁺/CLL-1⁺ AML cells are able to engraft and generate CLL-1⁺ blasts after transplantation in Non-Obese Diabetic/ Severe Combined ImmunoDeficient (NOD/SCID) mice [24].

CD96 is a transmembrane protein belonging to the Ig superfamily. The extracellular domain of CD96 has three Ig-like regions and a stalk-like region which can be heavily O-glycosylated. The only ligand for CD96 identified thus far is CD155, also known as polio virus receptor [26]. On normal cells CD96 protein expression appears to be restricted to T and NK cells, while its expression was also detected on T cell -Acute Lymphoid Leukemia (ALL) cells and AML cells [27;28]. Recently, Hosen and colleagues identified that compared with normal HSCs in AML-LSCs CD96 mRNA levels are at least 200-fold increased [28]. These mRNA expression data suggests that CD96 protein may be selectively over-expressed on AML-LSCs. In line with this finding, they demonstrated that CD96⁺ AML cells have a significantly higher engraftment potential than CD96⁻ AML cells [28]. Therefore, the development of antibody-based or CD96 ligand-based therapeutics may have potential for the targeted elimination of CD96⁺ AML-LSCs.

Targeting LSCs with naked antibodies

Indications for the efficacy of targeting CD123⁺ AML-LSCs with naked antibodies is demonstrated by a preclinical study using the CD123 neutralizing antibody, mAb7G3 [29]. Treatment with mAb7G3 inhibited engraftment and homing of human CD123⁺ AML-LSCs when injected in NOD/SCID mice. This effect of mAb7G3 was restricted to AML cells, since there was only a minor effect towards normal bone marrow (BM) cells. Furthermore, mAb7G3 treatment of NOD/SCID mice with pre-established AML reduced both the disease burden and the ability to transplant AML in secondary recipient mice (Lock R, et al. ASH Annual Meeting Abstracts 2007;110:161).

Recently, *in vitro* treatment with naked anti-CLL-1 mAbs in the presence of serum resulted in efficient Complement Dependent Cytotoxicity (CDC) towards various CLL-1⁺ AML cell lines. In line with these findings, *ex vivo* treatment of primary AML blasts with naked anti-CLL1 mAbs resulted in CDC in all samples tested. Furthermore, in a xenograft model one particular anti-CLL-1 mAb reduced the growth of established tumors by 38% (Korver W, et al. ASH Annual Meeting Abstracts 2008;112:4003). Together, these data indicate that targeting both CD123 and CLL-1 by naked antibodies might be a promising approach for the selective elimination of LSCs.

Targeting LSCs with immunoconjugates

The expression of CD33 in AML has resulted in the development of several CD33-targeted experimental therapeutics [30;31] and the clinically applied anti-AML immunotoxin Gemtuzumab Ozogamicin (GO). GO consists of an anti-CD33 mAb that is chemically

coupled to a derivative of the calicheamicin-toxin. After CD33-selective binding, GO internalizes and ends up in the acidic milieu of lysosomes where the calicheamicin moiety is hydrolytically released. Subsequently, calicheamicin translocates to the nucleus and intercalates with the DNA, causing site-specific double-strand breaks resulting in apoptotic cell death [32]. In the United States, GO was approved by the FDA for use in patients over the age of 60 with relapsed AML who are not considered candidates for standard chemotherapy [33]. In this subset of patients, GO treatment resulted in a 26% remission rate, albeit with sometimes considerable toxicity [34]. Recently, we have demonstrated that the activity of GO is strongly enhanced when treatment is combined with the epigenetic modifying drug valproic acid (VPA), a histone deacetylase inhibitor (HDACi). The co-treatment of AML cells with GO and VPA facilitated the intercalation of calicheamicin with the DNA, followed by synergistic apoptosis induction [35].

It has been reported that *in vitro* treatment of AML-LSCs with GO results in a median decrease in cellular viability with 46%, while normal HSCs were largely unaffected (Jawad M et al. ASH Annual Meeting Abstracts 2007;110:650). Therefore, it would be interesting to assess whether the anti-LSCs activity of GO can be further enhanced when treatment is combined with VPA.

One potential resistance mechanism of AML-LSCs towards treatment with GO involves the increased expression and/or activity of drug efflux pumps. In this respect, it appears that P-glycoprotein (Pgp) expression confers resistance to GO and as such is associated with a worse clinical response [36]. Therefore, combination treatment of GO and Pgp inhibitors may enhance the therapeutic effect of GO towards Pgp-expressing LSCs.

Alternatively, efflux pump-related resistance in AML-LSCs may be overcome using an anti-CD33 antibody conjugated to cytolytic radionuclides like ^{111}In . *In vitro* experiments with an ^{111}In -conjugated anti-CD33 mAb resulted in potent apoptosis in both chemo-resistant CD33⁺, HL-60 AML cells and in primary AML cells. Indeed, efflux pump-related resistance was overcome by ^{111}In since these AML cells expressed several drug efflux pumps including Pgp, Breast Cancer Resistance Protein 1 and Multidrug Resistance Protein 1 [37].

Currently, the efficacy of various CD123-targeted immunoconjugates is under investigation. In one particular approach CD123 is targeted by a fusion protein comprising a CD123-specific single chain fragment of the variable regions (scFv) antibody fragment that is genetically fused to a fragment of pseudomonas exotoxin A (PE38). This fusion protein, designated 26292(Fv)-PE38-KDEL, potentially induced apoptosis in a variety of CD123⁺ leukemia cell lines [38]. It remains to be investigated whether 26292(Fv)-PE38-KDEL can also eliminate CD123⁺ LSCs.

Targeting LSCs with ligand-based approaches

The cytokine IL3 has been used in fusion protein format to selectively deliver diphtheria toxin (DT) to CD123⁺ AML cells [39]. This IL3-DT fusion protein also selectively eliminated primitive AML progenitors, while normal progenitors were not affected [40]. Importantly, the IL3-DT fusion protein inhibited engraftment of human CD123⁺ AML-LSC cells when injected in NOD/SCID mice [40;41]. Recently, it was demonstrated that the pro-apoptotic activity of IL3-DT is enhanced when treatment is combined with Tumor necrosis factor

Related Apoptosis Inducing Ligand (TRAIL) [42]. Currently, a phase I clinical trial with this IL3-DT fusion protein for a subset of AML and MDS patients is initiated [41].




Figure 2. Antibody- and ligand-based approaches for the elimination of LSCs. The cell surface molecules CD96, CD123 (IL3Rα), CLL-1, CD33 and TRAIL-R1/R2 may be exploited for the selective elimination of LSCs. CD123 targeted strategies include mAb7G3, a fusion protein comprising an CD123-specific antibody fragment linked to pseudomonas endotoxin A (26292(Fv)-PE38-KDEL) and an IL3:diphtheria toxin (IL-3:DT) fusion protein. CLL-1 has been targeted with naked antibodies resulting in Complement Dependent Cytotoxicity. CD33 has been targeted with several experimental therapeutics, including radionuclides, a scFvCD33:sTRAIL fusion protein and the clinically applied immunotoxin Gemtuzumab Ozogamicin (GO). After CD33-selective binding, GO internalizes and ends up in the lysosomes. In the acidic milieu of the lysosomes, the calicheamicin moiety is hydrolytically released. Next, calicheamicin translocates to the nucleus and intercalates with the DNA, resulting in apoptotic cell death. Therapeutic strategies targeting the death-receptors, TRAIL-R1/R2, include activating anti-TRAIL-R1 and anti-TRAIL-R2 antibodies, recombinant preparations of soluble TRAIL (sTRAIL) and scFv:sTRAIL fusion proteins. Activation of TRAIL-R1/-R2 receptors results in the recruitment of the adaptor molecule FADD and caspase-8 to the death domains of these receptors. Assembly of this so-called death-inducing signaling complex (DISC) leads to sequential activation of caspase-8 and caspase-3, ultimately resulting in DNA damage and apoptosis. In some cell types, caspase-8 can cleave Bid (tBid) and thereby activate a mitochondrial amplification loop. This loop is characterized by cytochrome C release from the mitochondria and subsequent the sequential activation of caspase-9 and caspase-3.

Activation of apoptosis inducing receptors

The agonistic TRAIL death receptors, TRAIL-R1 and TRAIL-R2 (also known as DR4 and DR5, respectively) belong to the Tumor Necrosis Factor receptor family of death receptors. Ligation of these death receptors by TRAIL activates the extrinsic apoptotic pathway (reviewed in [43]). Hereupon, the adaptor protein FAS associated death domain (FADD) and the initiator caspase-8 are recruited to the intracellular death domains of these receptors. Assembly of this so-called death-inducing signaling complex (DISC)

leads to sequential activation of initiator and effector caspases and, ultimately, results in apoptotic cell death. In some cell types, the death-receptor pathway relies on an additional mitochondrial amplification loop that is activated by caspase-8-mediated cleavage of the BH3-only interacting domain death agonist (Bid) to a truncated form. Next, truncated BID (tBid) activates the mitochondrial pathway which involves cytochrome C and caspase-9. Intriguingly, apoptosis resulting from ligation of agonistic TRAIL-Rs appears to be restricted to malignant cells only, since normal cells appear to be resistant to TRAIL-R ligation. The mode of action of this tumor-selectivity of TRAIL-Rs ligation is still enigmatic.

Currently, activating anti-TRAIL-R1 and anti-TRAIL-R2 antibodies, as well as recombinant preparations of soluble TRAIL (sTRAIL) are evaluated in clinical trials [44-47]. In our laboratory, we have developed a series of fusion proteins, consisting of a tumor-selective scFv antibody fragment genetically fused to sTRAIL. These scFv:sTRAIL fusion proteins are designed to enhance tumor-selective cell surface delivery of sTRAIL (reviewed in [48]). Pre-clinically, these scFv:sTRAIL fusion proteins demonstrate potent and target antigen-restricted apoptosis induction. To this end, we have exploited both antigens expressed on solid tumors including Epidermal Growth Factor Receptor [49-51] and Epithelial Cell Adhesion Molecule [52;53], as well as antigens selectively expressed on leukemia cells including CD7 [54], CD19 [55] and CD33 [30].

Very recently, the expression of both TRAIL-R1 and TRAIL-R2 has been reported on LSCs. Comparing the gene expression profiles of AML-LSCs and HSCs, Majeti and colleagues demonstrated an increased expression of TRAIL-R1 and TRAIL-R2 in LSCs [56]. Furthermore, Yong et al. demonstrated that in CML non-cycling CD34⁺ cells have a significant higher expression of TRAIL-R1 and TRAIL-R2 than cycling CD34⁺ CML cells [57]. In line with this finding, sTRAIL significantly inhibited the clonogenic capacity of CD34⁺ AML cells without affecting the clonogenic capacity of normal CD34⁺ cells [58]. Together, these studies indicate that TRAIL-R1 and TRAIL-R2 receptors might be suitable candidates for the selective elimination of LSCs by targeted apoptosis induction. At present, we are assessing the efficacy of CD33- and CLL-1-selective scFv:sTRAIL fusion proteins for targeted apoptosis induction in AML-LSCs.

DISRUPTION OF THE LSCS-NICHE INTERACTION

In 1978, Schofield proposed that HSCs receive regulatory factors and signals within a specialized niche [59]. Currently, there are indications that these specialized niches are located in the bone marrow (BM), near the endosteum [60]. The endosteum is the inner surface located at the bone-BM interface and consisting mainly of osteoblasts and osteoclasts. More recent evidence indicates that also vascular and perivascular cells appear to contribute to these niche interactions as well (see review [61]). Signals within the HSCs niche regulate the normal survival, proliferation and differentiation of the HSCs. Recently, it was demonstrated that in leukemia the micromilieu in the HSCs niche appears to be disturbed by the presence of Stem Cell Factor that is locally secreted by leukemia cells [62]. Furthermore, it has been demonstrated that intravenously injected AML-LSCs home to, engraft and subsequently reside in the endosteal region [63]. Such cellular tropism of AML-LSCs for a dedicated niche typically involves the coordinated signaling by various chemokines and adhesion molecules. Neutralization or inhibition of these signals may

be exploited for the selective elimination of LSCs. Below, a selection of recent studies is reviewed that collectively indicate that disruption of the LSCs-niche interactions may indeed form a therapeutic option for the elimination of LSCs (summarized in Fig. 3).

Targeting the Stroma-derived factor-1(SDF-1)/CXCR4 axis

Previously, it has been demonstrated that the chemokine SDF-1 (CXCL12) as expressed by reticular stromal cells in the BM is pivotal for homing of HSCs to their niche [64]. SDF-1 binds to the transmembrane receptor CXCR4 which is expressed on HSCs [64;65], leukemic blasts and AML-LSCs [66]. Of note, a high expression of CXCR4 in AML correlates with a negative prognosis [67;68]. The pivotal role of CXCR4 in retention of HSCs in the niche is clearly demonstrated by the mobilization of HSCs by the CXCR4 antagonist AMD3100 [69].

The interaction of leukemia cells with cell adhesion molecules in the niche typically reduces their sensitivity to cytotoxic therapies, a process known as cell adhesion-mediated drug resistance (CAM-DR). Inhibition of CXCR4 by the specific inhibitors AMD3100 and AMD3465 sensitized AML cells for cytotoxic therapies [70-72]. Most likely, treatment with the CXCR4 inhibitors mobilizes AML cells, thereby reducing CAM-DR and sensitizing the AML cells for cytotoxic therapy.

Tavor and colleagues demonstrated that the anti-CXCR4 mAb12G5 significantly inhibits homing of AML cells to the BM of recipient mice. Notably, treatment of mice with established AML with mAb12G5 reduced the leukemic burden in the BM, blood and spleen. Importantly, similar treatment did not affect the levels of engrafted HPCs [66].

Additionally, lowering available SDF-1 levels in leukemia might offer a therapeutic strategy. Recently, the medicinal herb-derived compound berberine was shown to inhibit SDF-1 production by stromal cells, thereby impairing the migration of AML blasts and AML-LSCs *in vitro* [73]. Together these results indicate the importance of a properly functioning SDF-1/CXCR4 axis for homing and retention of AML blasts and AML-LSCs. Therefore, the SDF-1/CXCR4 axis appears to be a suitable target candidate for the selective elimination of AML-LSCs.

Targeting Very Late Antigen 4 (VLA-4)

The β 1-integrin, VLA-4, has a vital role in the retention of HPCs in the BM niche by immobilizing these cells [74]. The interaction of VLA-4 on AML cells with fibronectin on stromal cells initiates the phosphatidylinositol-3-OH kinase (PI3K) pro-survival pathway. Activation of the PI3K pathway in AML cells contributes to CAM-DR, which might explain the observation that VLA-4 expression on AML cells is correlated with a negative prognosis for AML patients. In line with this, Matsunaga and colleagues demonstrated that inhibition of VLA-4 interaction with fibronectin sensitized AML cells to cytosine arabinoside (Ara-C). This was demonstrated in a mice model for MRD in which they applied VLA-4 neutralizing antibodies to prevent the interaction with fibronectin [75]. Using the same MRD model, they recently demonstrated that a fibronectin-based synthetic peptide, designated FNIII14 [76], also sensitized AML cells to Ara-C [77]. Both approaches may be of considerable interest for disruption of the LSCs-niche interaction.

Targeting CD44

CD44 is a transmembrane receptor capable of binding several ligands, including hyaluronic acid which is abundantly expressed in the endosteal region [78]. On HSCs, CD44 serves as a homing receptor by interacting with E-selectin and L-selectin expressed in the BM [79]. Since CD44 is expressed on both CML-LSCs and AML-LSCs [80;81], neutralization of CD44-binding might impair the homing and retention of LSCs in the BM.

In CML, about 95% of the cases is characterized by the presence of the oncogenic fusion protein Bcr-Abl, a constitutively active mutant tyrosine kinase that causes uncontrolled cellular proliferation. The Bcr-Abl inhibitor Imatinib Mesylate induces remissions in the majority of CML patients. However, Imatinib appears to be unable to eradicate quiescent CML-LSCs [82], which might explain why the majority of CML patients relapse upon treatment discontinuation. Therefore, novel therapeutics for the selective elimination of Bcr-Abl⁺ CML-LSCs are needed.




Figure 3. Disruption of the LSC-niche interaction. The LSC-niche interaction can be disrupted by targeting VLA-4, CXCR4, CD44 and Rac1. Binding of LSCs-expressed VLA-4 to fibronectin in the BM initiates the PI3K pro-survival pathway and cell adhesion-mediated drug resistance (CAM-DR). The VLA-4-fibronectin interaction has been disrupted by mAbs and the synthetic peptide FNIII14. CXCR4 has been targeted with mAb12G5 and the specific inhibitors AMD3100 and AMD3465. Furthermore, the concentration of the CXCR4 ligand, SDF-1, can be lowered using berberine. The interaction of CD44 with the niche has been disrupted by mAbH90 and mAbIM7. The compound, NSC23766, has been utilized to specifically inhibit Rac1.

In a series of elegant experiments, Krause and colleagues demonstrated that CD44 expressed on Bcr-Abl⁺ CML-LSCs is essential for homing of these cells in recipient mice. Thereto, they performed a competitive homing assay in which the relative contribution of Bcr-Abl-transduced CD44⁺ BM cells and Bcr-Abl-transduced CD44⁻ BM cells in the development of CML was analyzed. Both cells types were mixed and injected intravenously

in irradiated recipient mice. After CML development, the animals were sacrificed and the relative contribution of the two distinct cell populations was analyzed. This revealed that Bcr-Abl-transduced CD44⁺ BM cells contributed 5 times more to the development of CML than Bcr-Abl-transduced CD44⁻ BM cells. CML development by the two different cell types was similar when injected intrafemorally. These results indicate that CD44 is necessary for homing of intravenously injected Bcr-Abl transduced cells to the BM. Importantly, anti-CD44 antibody treatment significantly prolonged survival of NOD/SCID recipient mice upon transplantation with CML-LSCs [80].

Simultaneously, the group of John Dick obtained similar results by targeting CD44 in AML. Treatment with the CD44-activating mAbH90 inhibited AML-LSCs homing to the BM niche and altered the fate of the AML-LSCs resulting in an efficient and selective elimination of the AML-LSCs [81]. Together, these studies clearly indicate a rationale for targeting CD44 to selectively eliminate CML-LSCs and AML-LSCs. Therefore, an evaluation of the efficacy of targeting CD44 in both CML and AML patients seems warranted.

Targeting Rac

The Rac subfamily of the Rho-family of GTP-ases is implicated in cell adhesion and motility. The family consists of three members Rac1, Rac2 and Rac3. Rac1 is ubiquitously expressed [83], while Rac2 expression is limited to the hematopoietic system [84] and Rac3 is selectively expressed in the brain [85]. Rac1^{-/-} HSCs and progenitors are unable to reconstitute hematopoiesis in irradiated recipient mice upon transplantation [86]. Moreover, it has been shown that Rac is overexpressed on CD34⁺ cells derived from AML patients. Rac appears to contribute to an enhanced migration and adhesion capacity of CD34⁺ cells to BM stromal cells.

Gao and colleagues [87] generated a Rac1 inhibitor, designated NSC23766, that significantly delayed leukemia development in a murine tumor model [88] and strongly reduced the clonal expansion of CD34⁺ AML cells [89]. Together, these promising data warrants further pre-clinical assessment of Rac1 inhibition for the elimination of LSCs.

TARGETING ABERRANT MOLECULAR PATHWAYS

Both, HSCs and LSCs rely on similar molecular pathways for their maintenance and self-renewal. However, in LSCs these pathways are often aberrantly regulated. LSCs appear to be particularly dependent on these deregulated pathways to maintain their malignant phenotype. This strong dependence indicates that therapeutic intervention in these pathways might be exploitable for the selective elimination of LSCs. Thus far, several of these deregulated pathways have been identified (e.g. the Hedgehog and Wnt pathways). The selective targeting of these pathways may yield promising anti-LSCs activity.

Compounds such as arsenic trioxide and parthenolide have been assessed for their capacity to induce apoptosis in LSCs. However, the pathways in which these compounds intervene are currently not completely understood. Therefore, further delineation of their specific molecular targets may contribute to development of more and rationally designed LSCs targeted therapeutics. Below, we provide a selection of aberrantly regulated pathways present in LSCs and a selection of chemical compounds that may have potential for the elimination of LSCs (summarized in Fig. 4).

The Hedgehog (Hh) pathway

The Hh pathway is involved in HSCs self-renewal [90] and is initiated by binding of one of the three Hh ligands, Sonic Hedgehog (Shh), Indian Hedgehog (Ihh) or desert Hedgehog (Dhh) to the transmembrane receptor Patched. Upon binding of Hh to Patched the inhibiting effect of Patched on the transmembrane receptor Smoothened (Smo) is relieved (reviewed in [91]). Next, Smo can activate Hh targets genes via the Gli family of transcription factors. Amongst these Hh targets genes are CyclinD/E [92] and B-cell-specific Moloney murine leukemia virus insertion site 1 (Bmi-1) [93] which are involved in cell proliferation and self-renewal, respectively. The Hh pathway can be inhibited by the chemical compound cyclopamine, which stabilizes Smo in its inactive form [94]. In a very recent study conducted by Zhao and colleagues the role of Smo in Bcr-Abl⁺ CML-LSCs was analyzed [95]. Thereto, Bcr-Abl-transduced normal HPCs and Bcr-Abl-transduced Smo^{-/-} HPCs were transplanted in irradiated recipient mice after which the development of CML was compared. More than 90% of the mice transplanted with Bcr-Abl-transduced normal HPCs developed CML. In contrast, only 47% of the mice developed CML when transplanted with Bcr-Abl-transduced Smo^{-/-} HPCs. When the Bcr-Abl-transduced HSCs were transplanted in mice, all recipient mice died from CML within four weeks. However, when these transplanted mice were treated with cyclopamine, 60% of the animals were still alive after 7 weeks. Furthermore, CML cells freshly isolated from cyclopamine-treated mice were unable to effectively repopulate secondary recipients. This corroborates with the strong reduction (up to 14-fold) of CML-LSCs in cyclopamine-treated mice. Additionally, cyclopamine treatment of human CML cell lines and primary CML samples reduced their clonogenic capacity. Importantly, the authors indicate that Smo inhibition by cyclopamine may also impaired the propagation of Imatinib-resistant human CML cells in xenografts [95].

The Wnt pathway

Wnt is implicated in HSCs self-renewal and proliferation [96] and is overexpressed in many different types of human cancers [97]. Thus far, three distinct Wnt signaling pathways have been described of which the canonical pathway is most prominently involved in leukemia. The key mediator of the canonical pathway is β -catenin. Normally, β -catenin is degraded rapidly after synthesis by the multiprotein destruction complex (MDC), which tags β -catenin for proteasomal degradation. The MDC consists of axin, adenomatous polyposis coli (APC) and glycogen-synthase kinase 3 β (GSK-3 β). Upon binding of Wnt to its receptors Frizzled and LRP5/6, the MDC becomes destabilized and therefore unable to tag β -catenin for degradation. Consequently, β -catenin accumulates and is able to translocate to the nucleus. In the nucleus β -catenin complexes with T-cell factor (TCF), B cell lymphoma 9 (Bcl-9) and pygo to induce TCF target genes expression [98], including the oncogenes C-MYC [99], Cyclin-D1 [100] and the cell surface antigen CD44 [101]. In AML, several oncogenic fusion proteins, such as AML-1/eight twenty one (AML1-ETO), promyelocytic leukemia / retinoic acid receptor alpha (PML/RAR α) and promyelocytic leukemia zinc finger (PLZF/RAR α) [102] can activate the Wnt signaling pathway. Furthermore, evidence is emerging that indicate an aberrant Wnt signaling pathway in LSCs [56] (extensively reviewed in ref [103]). In this respect, it has been shown that β -catenin plays a pivotal role in self-renewal of both HSCs as well as CML-LSCs [104]. Furthermore, very recently β -catenin has been implicated in the survival of Imatinib

resistant CML-LSCs in Bcr-Abl induced CML [105]. Therefore, modulation of Wnt signaling in the LSCs may contribute to their selective elimination.

There are several possibilities to inhibit the Wnt signaling pathway. Wnt signaling can be inhibited by extracellular proteins (reviewed in [106]) which scavenge Wnt and thereby prevent Wnt from binding to its receptors. These Wnt binding proteins include secreted Frizzled-related proteins (sFRP), Wnt inhibitory factor 1 (wif-1) and Cerberus. Moreover, members of the Dickkopf (DKK) family can directly bind the Wnt receptor and thereby prevent Wnt-induced signaling. In this respect, very recently it appeared that the anti-proliferative effects of human mesenchymal stem cells towards CML cells is at least partly attributable to secretion of DKK-1 [107]. Additionally, also very recently two natural compounds, CGP049090 and PKF 115-584 [108], have been identified that disrupt the TCF/ β -catenin complex. In AML these compounds possess potent tumoricidal activity [109]. It would be worthwhile to assess the possibilities of inhibition of the Wnt signaling pathway for the elimination of LSCs.

Aldehyde dehydrogenases (ALDH) enzymes

ALDH are a group of cytosolic enzymes that catalyzes the conversion of intracellular aldehydes. The ALDH superfamily consists of seventeen members [110] of which two members, cytosolic aldehyde dehydrogenase class-1A1 (ALDH1A1) and class-3A1 (ALDH3A1), have been implicated in tumor drug resistance against oxazaphosphorines (e.g. cyclophosphamide) [111]. ALDH enzyme activity, including the activity of ALDH1A1 and ALDH3A1, can be assessed using activatable fluorescent substrates. ALDH enzyme activity measurements enables the identification and isolation of HSCs [112], breast cancer stem cells [113], liver cancer stem cells [114] and AML-LSCs [115]. Manipulation of ALDH1A1 and ALDH3A1 expression or activity may be used to sensitize AML-LSCs to chemotherapy, especially to oxazaphosphorines. In this respect, the ALDH1 inhibitor, diethylamino-benzaldehyde (DEAB) might be utilized for sensitizing AML-LSCs to treatment with chemotherapeutic agents. Unfortunately, DEAB appeared to be very instable *in vivo* [116]. Therefore, the development of novel DEAB-analogues with enhanced stability is warranted. All-trans Retinoic Acid (ATRA), an established clinically applied anti-leukemia agent, appears to downregulate both ALDH1A1 and ALDH3A1 which may be beneficial for the selective elimination of LSCs [111]. In this respect, treatment of Acute Promyelocytic Leukemia (APL) and AML cells with ATRA resulted in differentiation of tumor cells and stimulation of cell cycling [117]. Therefore, we suggest that the combination treatment of oxazaphosphorines and ATRA may have potential for the elimination of LSCs.

Phosphatase and tensin homologue (PTEN)

PTEN is the second most commonly mutated gene in human malignancies, including hematopoietic malignancies [118;119]. Normally, PTEN inhibits cell proliferation and cell survival by negatively regulating the PI3K pathway [120]. In this respect, Yilmaz and colleagues recently compared the effect of PTEN inactivation in HSCs and LSCs [121]. Conditional deletion of PTEN resulted in the development of myeloproliferative disorder in 17 out of 19 mice which in the majority culminated in leukemia. In the HSCs, deletion of PTEN was characterized by an elevated cell cycle progression and an impaired replenishment of the HSCs pool, thus finally resulting in a depletion of HSCs. Apparently, PTEN deletion facilitates leukemogenesis while it depletes HSCs. These effects of PTEN

deletion appear to be mediated by mammalian target of rapamycin (mTOR), since treatment with the mTOR inhibitor rapamycin reversed the effects. Therefore, rapamycin treatment seems to be a promising approach for the selective elimination of LSCs while sparing HSCs.

However, Ito and colleagues recently reported on a contradictory effect of rapamycin [122]. They investigated the role of the tumor-suppressor promyelocytic leukemia protein (PML) in CML. Their results demonstrated that PML-deficient CML-LSCs are less quiescent than wild-type CML-LSCs, resulting in LSCs exhaustion. Furthermore, deletion of PML in HSCs also results in exhaustion and consequently impairment in long-term hematopoietic reconstitution. Since PML deficiency in both HSCs and LSCs is characterized by an increased mTOR activity they assessed the effect of rapamycin treatment. Here, rapamycin rescued the typical PML-deficient phenotype, thus exhaustion of the CML-LSCs and HSCs was prevented. Together, these seemingly contradictory results of mTOR inhibition in the elimination of LSCs indicate that further delineation of the role of mTOR in LSCs is needed.

Of note, since arsenic trioxide is known to decrease PML levels, Ito and colleagues continued with assessing arsenic trioxide for the elimination of CML-LSCs [122]. Treatment with arsenic trioxide resulted in the exit of the CML-LSCs from quiescence and therefore the CML-LSCs might become exhausted and more vulnerable to chemotherapeutics. Importantly, this cell cycle entry appeared to be more profound in LSCs than in HSCs, thereby generating a possible therapeutic window for the treatment of CML-LSCs by arsenic trioxide.

4-benzyl, 2-methyl, 1,2,4-thiadiazolidine, 3,5 dione (TDZD-8)

TDZD-8 was designed as a non-ATP competitive inhibitor of GSK-3 β for the treatment of Alzheimer's disease [123]. However, Guzman and colleagues uncovered potent tumoricidal activity of TDZD-8 towards leukemia cells. In a series of experiments they demonstrated that TDZD-8 exhibit strong anti-leukemia effects towards AML, CLL, ALL and blast crisis CML cells, without apparent toxic effects towards normal BM cells. Since a panel of other GSK-3 β inhibitors did not possess any comparable activity, the anti-leukemia effect is probably not governed by the inhibition of GSK-3 β . This notion is supported by the observation that β -catenin, of which the degradation is in part mediated by GSK-3 β , is elevated in CML-LSCs. Moreover, very recently GSK-3 β missplicing was observed in CML-LSCs and not in HSCs [124], together indicating that GSK-3 β is dispensable for leukemogenesis.

Importantly, TDZD-8 induced apoptosis in AML-LSCs and CML-LSCs without affecting HSCs [125]. In line with this, TDZD-8 specifically reduced the engraftment potential of AML-LSCs and not of HSCs. TDZD-8 is a highly hydrophobic agent which may result in a rapid loss of membrane integrity in leukemic cells by TDZD-8 insertions in the cellular membrane. The absence of this effect in HSCs might indicate that there is crucial difference in the membrane composition of LSCs and HSCs [125], which might open novel avenues for the selective elimination of LSCs.

Parthenolide (PTL)

PTL is a compound, derived from the plant feverfew, which eliminates cancer cells via reactive oxygen species and the inhibition of NF- κ B [126-128]. Guzman and colleagues

analyzed the effects of PTL and Ara-C on AML blasts and AML-LSCs. Apoptosis induction in the AML blasts by Ara-C was more pronounced than by PTL. However, PTL potently induced apoptosis in AML-LSCs, whereas Ara-C exhibited only minimal apoptotic activity. Importantly, the PTL concentration necessary for significant AML-LSCs elimination had only a modest effect on HSCs [129]. Moreover, PTL treatment significantly reduced the engraftment potential of AML-LSCs, while PTL treatment did not affect the engraftment of HSCs [129].

Despite the favorable characteristics, the clinical application of PTL is probably hampered by its poor water solubility. Therefore, Guzman and colleagues generated a series of parthenolide analogues aiming to identify an active compound with improved pharmacological properties [130]. Recently, they identified a novel compound, designated dimethylamino analog of parthenolide (DMAPT), which exhibits similar activity as PTL. However, when formulated as a fumarate salt, DMAPT had 1000-fold enhanced water solubility, resulting in a 70% bioavailability of orally applied DMAPT [130]. In contrast to PTL, oral administration of DMAPT did result in plasma concentrations sufficient to exert an anti-leukemia effect in both rodent and canine models.

EPIGENETIC REPROGRAMMING OF LSCS

The step-wise conversion of a normal cell to a leukemic cell typically involves the activation of oncogenes and anti-apoptotic genes and the inactivation of tumor suppressor and pro-apoptotic genes [131]. Frequently, gene expression in leukemic cells is altered by epigenetic reprogramming (reviewed in [132]).

Epigenetic reprogramming is characterized by various distinct chromatin modifications. Chromatin consists of histone proteins and DNA. The central unit of the chromatin structure is an octamer of core histone proteins, referred to as the nucleosome. The distance between adjacent nucleosomes and thus the compaction of the chromatin in between, defines two types of chromatin. Euchromatin is the open type of chromatin with a relative low compaction which allows access of the transcription machinery to the DNA, while heterochromatin is the condensed form in which the transcription machinery can not access the DNA and reflects an inactive state. There are several types of modifications that can alter the accessibility of the transcription machinery to the DNA. The best studied are DNA and histone methylation which are associated with heterochromatin and histone acetylation which is associated with euchromatin.

Normal stem cells rely on Polycomb group (PcG) proteins to repress genes encoding transcription factors required for differentiation including the homeotic (HOX) genes [133]. It is postulated that methylation of promoter regions of those repressed genes could lock in stem cell phenotypes and initiate abnormal clonal expansion and thereby predispose to cancer [134;135]. Repression of genes by the PcG proteins is initiated by the Polycomb Repressor Complex 2 (PRC2) containing the histone methyltransferase EZH2, after which the Polycomb Repressor Complex 1 (PRC1) further maintains the repression. The PRC1 comprises amongst others Bmi-1, which is essential for self-renewal of HSCs [136]. Of note, in both AML and CML patients Bmi-1 is overexpressed compared to normal CD34+ BM cells. Patients with an elevated Bmi-1 expression have less chance of achieving complete remission, higher chance of relapse and reduced duration of survival [137;138]. Moreover, very recently Wang and colleagues have shown that the dysregulation of a

chromatin-binding plant homeodomain (PHD) finger can cause hematological malignancies [139]. Normally, PHD fingers respond to certain histone methylation states [140;141]. In AML however, chromosomal translocations are observed which results in the fusion of a PHD finger to a common leukemia fusion partner called NUP98 [142-144]. Wang and colleagues compared the transcriptome profile of BM progenitor cells transduced with such a fusion protein with control cells. This analysis indicated that in the cells transduced with this fusion protein several genes were upregulated including HOXA9 and MEIS1 [145]. Lessard and Sauvageau have shown that enforced expression of HOXA9/MEIS1




Figure 4. Targeting aberrantly regulated pathways and epigenetic reprogramming Aberrantly regulated Wnt and Hedgehog pathways have been identified in LSCs and therefore might be suitable candidates for therapeutic intervention. At the cell surface, binding of Wnt to LRP5/6 and Frizzled destabilizes the Multiprotein Destruction Complex (MDC), which in the absence of Wnt signaling facilitates the degradation of β -catenin. As a result of Wnt binding to its receptors, β -catenin accumulates and translocates to the nucleus where it complexes with TCF, Bcl-9 and pygo to induce the expression of several genes including c-myc, cyclin-D and CD44. Extracellularly, the Wnt signaling pathway can be inhibited by the scavengers sFRP, Wif1 and Cerberus and the DKK-family members which prevent Wnt binding to its receptor. Intracellularly, Wnt signaling can be inhibited by the compounds CGP049090 and PKF115-584, which disrupt the TCF/ β -catenin complex. The Hedgehog pathway is initiated by binding of a ligand (Shh, Ihh or Dhh) to the receptor Patched. Hereupon, Smoothened becomes activated and via the Gli family of transcription factors induces the expression of several target genes including CyclinD/E and Bmi-1. The Hedgehog pathway can be inhibited by cyclopamine which stabilizes Smoothened in its inactive form. Binding of Growth factors (GF) to Growth Factor Receptors (GFR) or binding of VLA-4 to fibronectin induces activation of PI3K. This pro-survival signaling pathway is characterized by the conversion of PIP2 to PIP3 and subsequent downstream signaling via several proteins, including mTOR. Normally, PTEN functions as an inhibitor of the PI3K pathway. Pharmacologically, both rapamycin and arsenic trioxide (via PML) can alter mTOR activation. The enzymes ALDH1A1 and ALDH3A1 have been implicated in Multi Drug Resistance (MDR) and can be inhibited by DEAB and ATRA. The anti-leukemia activity of parthenolide (PTL) is characterized by the inhibition of NF- κ B and the generation of ROS. The compound TDZD-8 can also inhibit NF- κ B activation and might influence LSCs membrane integrity. Frequently, gene expression in leukemic cells is altered by epigenetic reprogramming. Therefore, reprogramming the epigenetic status using decitabine or HDACi appears promising for the elimination of LSCs.

can induce leukemia, which is dependent on the expression of Bmi-1 [146], suggesting that Bmi-1 inhibitors may contribute to the elimination of LSCs.

However, also drugs with a more general epigenetic modification activity may have clinical relevance in attacking LSCs. This is evident from the clinical application of the demethylating agent decitabine, which is approved for the treatment of MDS. Decitabine inhibits DNA methyltransferases (DNMT) and therefore has a broad range of action by tipping the balance towards euchromatin. Intriguingly, decitabine exerts differential effects in LSCs and HSCs, which is probably due to differences in their epigenomic status. In LSCs treatment with decitabine results in differentiation, whereas decitabine treatment of HSCs results in enhanced self-renewal (Negrotto S et al. ASH Annual Meeting Abstracts 2008;112:201). Furthermore, it was demonstrated that decitabine treatment reactivated more genes in bladder cancer cells than in normal fibroblasts [147]. These studies indicate that general epigenetic modifications can exert differential effects in LSCs and HSCs, which might be utilized for the selective elimination of LSCs.

Other general epigenetic modulators are HDACi such as suberoylanilide hydroxamic acid (SAHA), 4-phenylbutyric acid (PBA) and VPA. HDACi can also tip the balance toward euchromatin and have shown to induce apoptosis in leukemia (see Fig. 4). In clinical trials VPA monotherapy was well tolerated, but showed disappointing anti-leukemia effects (reviewed in [148]). However, combination treatment of VPA with other anti-cancer agents showed significantly enhanced and sometimes synergistic anti-tumor effects [35;149-151].

MicroRNA's (miRs) are small, non-coding, single stranded RNA sequences that regulate gene expression by binding to mRNA molecules. Interestingly, upon co-treatment with decitabine and PBA, miRs appear to be differentially reactivated in cancer cells and normal cells [152]. Saito and colleagues have shown that the transcription of one particular miR, miR-127, was increased upon combination treatment of decitabine and PBA. Subsequently, they identified that a predicted target of miR-127, BCL6, was downregulated [152]. This indicates that not only epigenetically silenced protein-coding sequences, but also miRs can be reactivated upon treatment with demethylating agents and HDACi. This opens novel avenues for modifying protein levels in LSCs and therefore may be exploited to eliminate LSCs.

CONCLUSIONS AND PERSPECTIVES

The specific elimination of LSCs in hemato-oncology is a research area still in its infancy. Currently, researchers are at the brink of identifying biological processes imperative to LSCs biology. Elucidation of these LSCs-specific characteristics is of eminent importance for the rational design of novel LSCs targeted therapies. It is anticipated that the successful elimination of LSCs will have tremendous impact on leukemia therapy. Several strategies appear promising for the elimination of LSCs, including targeting LSC-selective cell surface antigens, disrupting LSCs-niche interactions, targeting aberrantly regulated pathways and epigenetic reprogramming of LSCs. Several studies utilizing these strategies have generated promising pre-clinical results. However, the vast majority of the data is obtained by xenografting LSCs in severely immunocompromised recipient mice. It is evident that the data obtained using these models can not be extrapolated directly to the clinical situation. Furthermore, it is likely that multifaceted therapy combining several

strategies will be the most effective approach. Therefore, rational designed combinatorial approaches and multi-modality targeting of LSCs are warranted. Subsequent, these combinatorial strategies have to prove to be safe and effective in clinical trials.

ACKNOWLEDGEMENTS

Supported by grants from the Dutch Cancer Society (2005-3358 and 2007-3784) to W.H.

REFERENCES

1. Jordan CT, Guzman ML, Noble M. Cancer stem cells *N.Engl.J.Med.* 2006;355:1253-1261
2. Krause DS and Van Etten RA. Right on target: eradicating leukemic stem cells *Trends Mol.Med.* 2007;13:470-481
3. Bonnet D and Dick JE. Human acute myeloid leukemia is organized as a hierarchy that originates from a primitive hematopoietic cell *Nat.Med.* 1997;3:730-737
4. Lapidot T, Sirard C, Vormoor J, Murdoch B, Hoang T, Caceres-Cortes J, Minden M, Paterson B, Caligiuri MA, Dick JE. A cell initiating human acute myeloid leukaemia after transplantation into SCID mice *Nature* 1994;367:645-648
5. Castor A, Nilsson L, Astrand-Grundstrom I, Buitenhuis M, Ramirez C, Anderson K, Strombeck B, Garwicz S, Bekassy AN, Schmiegelow K, Lausen B, Hokland P, Lehmann S, Juliusson G, Johansson B, Jacobsen SE. Distinct patterns of hematopoietic stem cell involvement in acute lymphoblastic leukemia *Nat.Med.* 2005;11:630-637
6. Misaghian N, Ligresti G, Steelman LS, et al. Targeting the leukemic stem cell: the Holy Grail of leukemia therapy *Leukemia* 2009;23:25-42
7. Dalerba P, Cho RW, Clarke MF. Cancer stem cells: models and concepts *Annu.Rev.Med.* 2007;58:267-284
8. Hope KJ, Jin L, Dick JE. Acute myeloid leukemia originates from a hierarchy of leukemic stem cell classes that differ in self-renewal capacity *Nat.Immunol.* 2004;5:738-743
9. Terpstra W, Ploemacher RE, Prins A, van Lom K, Pouwels K, Wognum AW, Wagemaker G, Lowenberg B, Wielenga JJ. Fluorouracil selectively spares acute myeloid leukemia cells with long-term growth abilities in immunodeficient mice and in culture *Blood* 1996;88:1944-1950
10. Dean M, Fojo T, Bates S. Tumour stem cells and drug resistance *Nat.Rev.Cancer* 2005;5:275-284
11. Lane SW, Scadden DT, Gilliland DG. The leukemic stem cell niche - current concepts and therapeutic opportunities *Blood* 2009;
12. Stel AJ, Ten Cate B, Jacobs S, Kok JW, Spierings DC, Dondorff M, Helfrich W, Kluijn-Nelemans HC, de Leij LF, Withoff S, Kroesen BJ. Fas receptor clustering and involvement of the death receptor pathway in rituximab-mediated apoptosis with concomitant sensitization of lymphoma B cells to fas-induced apoptosis *J.Immunol.* 2007;178:2287-2295
13. Byrd JC, Kitada S, Flinn IW, Aron JL, Pearson M, Lucas D, Reed JC. The mechanism of tumor cell clearance by rituximab in vivo in patients with B-cell chronic lymphocytic leukemia: evidence of caspase activation and apoptosis induction *Blood* 2002;99:1038-1043
14. Sato T, Laver JH, Ogawa M. Reversible expression of CD34 by murine hematopoietic stem cells *Blood* 1999;94:2548-2554
15. Freeman SD, Kelm S, Barber EK, Crocker PR. Characterization of CD33 as a new member of the sialoadhesin family of cellular interaction molecules *Blood* 1995;85:2005-2012
16. Wellhausen SR and Peiper SC. CD33: biochemical and biological characterization and evaluation of clinical relevance *J.Biol.Regul.Homeost.Agents* 2002;16:139-143
17. Tchilian EZ, Beverley PC, Young BD, Watt SM. Molecular cloning of two isoforms of the murine homolog of the myeloid CD33 antigen *Blood* 1994;83:3188-3198

18. Florian S, Sonneck K, Hauswirth AW, Krauth MT, Scherthaner GH, Sperr WR, Valent P. Detection of molecular targets on the surface of CD34+/CD38-- stem cells in various myeloid malignancies *Leuk.Lymphoma* 2006;47:207-222
19. Hauswirth AW, Florian S, Printz D et al. Expression of the target receptor CD33 in CD34+/CD38-/CD123+ AML stem cells *Eur.J.Clin.Invest* 2007;37:73-82
20. Jordan CT, Upchurch D, Szilvassy SJ, Guzman ML, Howard DS, Pettigrew AL, Meyerrose T, Rossi R, Grimes B, Rizzieri DA, Luger SM, Phillips GL. The interleukin-3 receptor alpha chain is a unique marker for human acute myelogenous leukemia stem cells *Leukemia* 2000;14:1777-1784
21. Moretti S, Lanza F, Dabusti M, Tieghi A, Campioni D, Dominici M, Castoldi GL. CD123 (interleukin 3 receptor alpha chain) *J.Biol.Regul.Homeost.Agents* 2001;15:98-100
22. Marshall AS, Willment JA, Lin HH, Williams DL, Gordon S, Brown GD. Identification and characterization of a novel human myeloid inhibitory C-type lectin-like receptor (MICL) that is predominantly expressed on granulocytes and monocytes *J.Biol.Chem.* 2004;279:14792-14802
23. Bakker AB, van den Oudenrijn S, Bakker AQ, Feller N, van Meijer M, Bia JA, Jongeneelen MA, Visser TJ, Bijl N, Geuijen CA, Marissen WE, Radosevic K, Throsby M, Schuurhuis GJ, Ossenkoppele GJ, de Kruif J, Goudsmit J, Kruisbeek AM. C-type lectin-like molecule-1: a novel myeloid cell surface marker associated with acute myeloid leukemia *Cancer Res.* 2004;64:8443-8450
24. van Rhenen A, van Dongen GA, Kelder A, Rombouts EJ, Feller N, Moshaver B, Stigter-van Walsum M, Zweegman S, Ossenkoppele GJ, Jan Schuurhuis G. The novel AML stem cell associated antigen CLL-1 aids in discrimination between normal and leukemic stem cells *Blood* 2007;110:2659-2666
25. Moshaver B, van Rhenen A, Kelder A, van der Pol M, Terwijn M, Bachas C, Westra AH, Ossenkoppele GJ, Zweegman S, Schuurhuis GJ. Identification of a small subpopulation of candidate leukemia-initiating cells in the side population of patients with acute myeloid leukemia *Stem Cells* 2008;26:3059-3067
26. Fuchs A, Cella M, Giurisato E, Shaw AS, Colonna M. Cutting edge: CD96 (tactile) promotes NK cell-target cell adhesion by interacting with the poliovirus receptor (CD155) *J.Immunol.* 2004;172:3994-3998
27. Gramatzki M, Ludwig WD, Burger R, Moos P, Rohwer P, Grunert C, Sender A, Kalden JR, Andreesen R, Henschke F, Moldenhauer G. Antibodies TC-12 ("unique") and TH-111 (CD96) characterize T-cell acute lymphoblastic leukemia and a subgroup of acute myeloid leukemia *Exp.Hematol.* 1998;26:1209-1214
28. Hosen N, Park CY, Tatsumi N, Oji Y, Sugiyama H, Gramatzki M, Krensky AM, Weissman IL. CD96 is a leukemic stem cell-specific marker in human acute myeloid leukemia *Proc.Natl.Acad.Sci.U.S.A* 2007;104:11008-11013
29. Sun Q, Woodcock JM, Rapoport A, Stomski FC, Korpelainen EI, Bagley CJ, Goodall GJ, Smith WB, Gamble JR, Vadas MA, Lopez AF. Monoclonal antibody 7G3 recognizes the N-terminal domain of the human interleukin-3 (IL-3) receptor alpha-chain and functions as a specific IL-3 receptor antagonist *Blood* 1996;87:83-92
30. ten Cate B, Bremer E, de Bruyn M, Bijma T, Samplonius D, Schwemmlin M, Huls G, Fey G, Helfrich W. A novel AML-selective TRAIL fusion protein that is superior to Gemtuzumab Ozogamicin in terms of in vitro selectivity, activity and stability *Leukemia* 2009;
31. Schwemmlin M, Peipp M, Barbin K, Saul D, Stockmeyer B, Repp R, Birkmann J, Oduncu F, Emmerich B, Fey GH. A CD33-specific single-chain immunotoxin mediates potent apoptosis of cultured human myeloid leukaemia cells *Br.J.Haematol.* 2006;133:141-151
32. van Der Velden VH, te Marvelde JG, Hoogeveen PG, Bernstein ID, Houtsmuller AB, Berger MS, van Dongen JJ. Targeting of the CD33-calicheamicin immunoconjugate Mylotarg (CMA-676) in acute myeloid leukemia: in vivo and in vitro saturation and internalization by leukemic and normal myeloid cells *Blood* 2001;97:3197-3204
33. Bross PF, Beitz J, Chen G, Chen XH, Duffy E, Kieffer L, Roy S, Sridhara R, Rahman A, Williams G, Pazdur R. Approval summary: gemtuzumab ozogamicin in relapsed acute myeloid leukemia *Clin.Cancer Res.* 2001;7:1490-1496
34. Larson RA, Sievers EL, Stadtmauer EA et al. Final report of the efficacy and safety of gemtuzumab ozogamicin (Mylotarg) in patients with CD33-positive acute myeloid leukemia in first recurrence *Cancer* 2005;104:1442-1452

35. ten Cate B, Samplonius DF, Bijma T, de Leij LF, Helfrich W, Bremer E. The histone deacetylase inhibitor valproic acid potentially augments gemtuzumab ozogamicin-induced apoptosis in acute myeloid leukemic cells *Leukemia* 2007;21:248-252
36. Walter RB, Gooley TA, van Der Velden VH, Loken MR, van Dongen JJ, Flowers DA, Bernstein ID, Appelbaum FR. CD33 expression and P-glycoprotein-mediated drug efflux inversely correlate and predict clinical outcome in patients with acute myeloid leukemia treated with gemtuzumab ozogamicin monotherapy *Blood* 2007;109:4168-4170
37. Kersemans V, Cornelissen B, Minden MD, Brandwein J, Reilly RM. Drug-resistant AML cells and primary AML specimens are killed by 111In-anti-CD33 monoclonal antibodies modified with nuclear localizing peptide sequences *J.Nucl.Med.* 2008;49:1546-1554
38. Du X, Ho M, Pastan I. New immunotoxins targeting CD123, a stem cell antigen on acute myeloid leukemia cells *J.Immunother.* 2007;30:607-613
39. Testa U, Riccioni R, Biffoni M, Diverio D, Lo-Coco F, Foa R, Peschle C, Frankel AE. Diphtheria toxin fused to variant human interleukin-3 induces cytotoxicity of blasts from patients with acute myeloid leukemia according to the level of interleukin-3 receptor expression *Blood* 2005;106:2527-2529
40. Feuring-Buske M, Frankel AE, Alexander RL et al. A diphtheria toxin-interleukin 3 fusion protein is cytotoxic to primitive acute myeloid leukemia progenitors but spares normal progenitors *Cancer Res.* 2002;62:1730-1736
41. Yalcintepe L, Frankel AE, Hogge DE. Expression of interleukin-3 receptor subunits on defined subpopulations of acute myeloid leukemia blasts predicts the cytotoxicity of diphtheria toxin interleukin-3 fusion protein against malignant progenitors that engraft in immunodeficient mice *Blood* 2006;108:3530-3537
42. Horita H, Frankel AE, Thorburn A. Acute-myeloid-leukemia-targeted toxins kill tumor cells by cell-type-specific mechanisms and synergize with TRAIL to allow manipulation of the extent and mechanism of tumor cell death *Leukemia* 2008;22:652-655
43. Ashkenazi A. Targeting the extrinsic apoptosis pathway in cancer *Cytokine Growth Factor Rev.* 2008;19:325-331
44. Tolcher AW, Mita M, Meropol NJ et al. Phase I pharmacokinetic and biologic correlative study of mapatumumab, a fully human monoclonal antibody with agonist activity to tumor necrosis factor-related apoptosis-inducing ligand receptor-1 *J.Clin.Oncol.* 2007;25:1390-1395
45. Hotte SJ, Hirte HW, Chen EX et al. A phase 1 study of mapatumumab (fully human monoclonal antibody to TRAIL-R1) in patients with advanced solid malignancies *Clin Cancer Res.* 2008;14:3450-3455
46. Plummer R, Attard G, Pacey S et al. Phase 1 and pharmacokinetic study of lexatumumab in patients with advanced cancers *Clin Cancer Res.* 2007;13:6187-6194
47. Ashkenazi A. Directing cancer cells to self-destruct with pro-apoptotic receptor agonists *Nat.Rev.Drug Discov.* 2008;7:1001-1012
48. Bremer E, de Bruyn M, Wajant H, Helfrich W. Targeted cancer immunotherapy using ligands of the tumor necrosis factor super-family *Curr.Drug Targets.* 2009;10:94-103
49. Bremer E, van Dam GM, de Bruyn M, van Riesen M, Dijkstra M, Kamps G, Helfrich W, Haisma H. Potent systemic anticancer activity of adenovirally expressed EGFR-selective TRAIL fusion protein *Mol.Ther.* 2008;16:1919-1926
50. Bremer E, Samplonius DF, van Genne L, Dijkstra MH, Kroesen BJ, de Leij LF, Helfrich W. Simultaneous inhibition of epidermal growth factor receptor (EGFR) signaling and enhanced activation of tumor necrosis factor-related apoptosis-inducing ligand (TRAIL) receptor-mediated apoptosis induction by an scFv:sTRAIL fusion protein with specificity for human EGFR *J Biol.Chem.* 2005;280:10025-10033
51. Bremer E, de Bruyn M, Samplonius DF, Bijma T, ten Cate B, de Leij LF, Helfrich W. Targeted delivery of a designed sTRAIL mutant results in superior apoptotic activity towards EGFR-positive tumor cells *J Mol.Med.* 2008;86:909-924
52. Bremer E, Samplonius D, Kroesen BJ, van Genne L, de Leij L, Helfrich W. Exceptionally potent anti-tumor bystander activity of an scFv:sTRAIL fusion protein with specificity for EGP2 toward target antigen-negative tumor cells *Neoplasia.* 2004;6:636-645

53. Bremer E, Kuijlen J, Samplonius D, Walczak H, de Leij L, Helfrich W. Target cell-restricted and -enhanced apoptosis induction by a scFv:sTRAIL fusion protein with specificity for the pancarcinoma-associated antigen EGP2 *Int.J Cancer* 2004;109:281-290
54. Bremer E, Samplonius DF, Peipp M, van Genne L, Kroesen BJ, Fey GH, Gramatzki M, de Leij LF, Helfrich W. Target cell-restricted apoptosis induction of acute leukemic T cells by a recombinant tumor necrosis factor-related apoptosis-inducing ligand fusion protein with specificity for human CD7 *Cancer Res.* 2005;65:3380-3388
55. Stieglmaier J, Bremer E, Kellner C, Liebig TM, ten Cate B, Peipp M, Schulze-Koops H, Pfeiffer M, Buhring HJ, Greil J, Oduncu F, Emmerich B, Fey GH, Helfrich W. Selective induction of apoptosis in leukemic B-lymphoid cells by a CD19-specific TRAIL fusion protein *Cancer Immunol.Immunother.* 2008;57:233-246
56. Majeti R, Becker MW, Tian Q, Lee TL, Yan X, Liu R, Chiang JH, Hood L, Clarke MF, Weissman IL. Dysregulated gene expression networks in human acute myelogenous leukemia stem cells *Proc.Natl.Acad.Sci.U.S.A* 2009;
57. Yong A, Keyvanfar K, Hensel N, Eniafe R, Savani B, Berg M, Lundqvist A, Adams S, Sloan E, Goldman J, Childs R, Barrett AJ. Primitive quiescent CD34+ cells in chronic myeloid leukemia are targeted by in vitro expanded natural killer cells, which are functionally enhanced by bortezomib *Blood* 2009;113:875-882
58. Tazzari PL, Tabellini G, Ricci F, Papa V, Bortul R, Chiarini F, Evangelisti C, Martinelli G, Bontadini A, Cocco L, McCubrey JA, Martelli AM. Synergistic proapoptotic activity of recombinant TRAIL plus the Akt inhibitor Perifosine in acute myelogenous leukemia cells *Cancer Res.* 2008;68:9394-9403
59. Schofield R. The relationship between the spleen colony-forming cell and the haemopoietic stem cell *Blood Cells* 1978;4:7-25
60. Nilsson SK, Johnston HM, Coverdale JA. Spatial localization of transplanted hemopoietic stem cells: inferences for the localization of stem cell niches *Blood* 2001;97:2293-2299
61. Kiel MJ and Morrison SJ. Uncertainty in the niches that maintain haematopoietic stem cells *Nat.Rev.Immunol.* 2008;8:290-301
62. Colmone A, Amorim M, Pontier AL, Wang S, Jablonski E, Sipkins DA. Leukemic cells create bone marrow niches that disrupt the behavior of normal hematopoietic progenitor cells *Science* 2008;322:1861-1865
63. Ishikawa F, Yoshida S, Saito Y, Hijikata A, Kitamura H, Tanaka S, Nakamura R, Tanaka T, Tomiyama H, Saito N, Fukata M, Miyamoto T, Lyons B, Ohshima K, Uchida N, Taniguchi S, Ohara O, Akashi K, Harada M, Shultz LD. Chemotherapy-resistant human AML stem cells home to and engraft within the bone-marrow endosteal region *Nat.Biotechnol.* 2007;25:1315-1321
64. Peled A, Petit I, Kollet O et al. Dependence of human stem cell engraftment and repopulation of NOD/SCID mice on CXCR4 *Science* 1999;283:845-848
65. Sugiyama T, Kohara H, Noda M, Nagasawa T. Maintenance of the hematopoietic stem cell pool by CXCL12-CXCR4 chemokine signaling in bone marrow stromal cell niches *Immunity.* 2006;25:977-988
66. Tavor S, Petit I, Porozov S, Avigdor A, Dar A, Leider-Trejo L, Shemtov N, Deutsch V, Naparstek E, Nagler A, Lapidot T. CXCR4 regulates migration and development of human acute myelogenous leukemia stem cells in transplanted NOD/SCID mice *Cancer Res.* 2004;64:2817-2824
67. Spoo AC, Lubbert M, Wierda WG, Burger JA. CXCR4 is a prognostic marker in acute myelogenous leukemia *Blood* 2007;109:786-791
68. Rombouts EJ, Pavic B, Lowenberg B, Ploemacher RE. Relation between CXCR-4 expression, Flt3 mutations, and unfavorable prognosis of adult acute myeloid leukemia *Blood* 2004;104:550-557
69. Broxmeyer HE, Orschell CM, Clapp DW, Hangoc G, Cooper S, Plett PA, Liles WC, Li X, Graham-Evans B, Campbell TB, Calandra G, Bridger G, Dale DC, Srour EF. Rapid mobilization of murine and human hematopoietic stem and progenitor cells with AMD3100, a CXCR4 antagonist *J.Exp.Med.* 2005;201:1307-1318
70. Nervi B, Ramirez P, Rettig MP, et al. Chemosensitization of acute myeloid leukemia (AML) following mobilization by the CXCR4 antagonist AMD3100 *Blood* 2009;113:6206-6214

71. Zeng Z, Shi YX, Samudio IJ, Wang RY, Ling X, Frolova O, Levis M, Rubin JB, Negrin RR, Estey EH, Konoplev S, Andreeff M, Konopleva M. Targeting the leukemia microenvironment by CXCR4 inhibition overcomes resistance to kinase inhibitors and chemotherapy in AML *Blood* 2009;113:6215-6224
 72. Burger JA and Peled A. CXCR4 antagonists: targeting the microenvironment in leukemia and other cancers *Leukemia* 2009;23:43-52
 73. Li H, Guo L, Jie S et al. Berberine inhibits SDF-1-induced AML cells and leukemic stem cells migration via regulation of SDF-1 level in bone marrow stromal cells *Biomed. Pharmacother.* 2008;62:573-578
 74. Papayannopoulou T, Priestley GV, Nakamoto B, Zafiropoulos V, Scott LM. Molecular pathways in bone marrow homing: dominant role of alpha(4)beta(1) over beta(2)-integrins and selectins *Blood* 2001;98:2403-2411
 75. Matsunaga T, Takemoto N, Sato T, Takimoto R, Tanaka I, Fujimi A, Akiyama T, Kuroda H, Kawano Y, Kobune M, Kato J, Hirayama Y, Sakamaki S, Kohda K, Miyake K, Niitsu Y. Interaction between leukemic-cell VLA-4 and stromal fibronectin is a decisive factor for minimal residual disease of acute myelogenous leukemia *Nat. Med.* 2003;9:1158-1165
 76. Kato R, Ishikawa T, Kamiya S, Oguma F, Ueki M, Goto S, Nakamura H, Katayama T, Fukai F. A new type of antimetastatic peptide derived from fibronectin *Clin. Cancer Res.* 2002;8:2455-2462
 77. Matsunaga T, Fukai F, Miura S, Nakane Y, Owaki T, Kodama H, Tanaka M, Nagaya T, Takimoto R, Takayama T, Niitsu Y. Combination therapy of an anticancer drug with the FNIII14 peptide of fibronectin effectively overcomes cell adhesion-mediated drug resistance of acute myelogenous leukemia *Leukemia* 2008;22:353-360
 78. Avigdor A, Goichberg P, Shvitiel S, Dar A, Peled A, Samira S, Kollet O, Herschkoviz R, Alon R, Hardan I, Ben Hur H, Naor D, Nagler A, Lapidot T. CD44 and hyaluronic acid cooperate with SDF-1 in the trafficking of human CD34+ stem/progenitor cells to bone marrow *Blood* 2004;103:2981-2989
 79. Sackstein R, Merzaban JS, Cain DW, Dagia NM, Spencer JA, Lin CP, Wohlgemuth R. Ex vivo glycan engineering of CD44 programs human multipotent mesenchymal stromal cell trafficking to bone *Nat. Med.* 2008;14:181-187
 80. Krause DS, Lazarides K, von Andrian UH, Van Etten RA. Requirement for CD44 in homing and engraftment of BCR-ABL-expressing leukemic stem cells *Nat. Med.* 2006;12:1175-1180
 81. Jin L, Hope KJ, Zhai Q, Smadja-Joffe F, Dick JE. Targeting of CD44 eradicates human acute myeloid leukemic stem cells *Nat. Med.* 2006;12:1167-1174
 82. Graham SM, Jorgensen HG, Allan E et al. Primitive, quiescent, Philadelphia-positive stem cells from patients with chronic myeloid leukemia are insensitive to STI571 in vitro *Blood* 2002;99:319-325
 83. Moll J, Sansig G, Fattori E, van der Putten H. The murine rac1 gene: cDNA cloning, tissue distribution and regulated expression of rac1 mRNA by disassembly of actin microfilaments *Oncogene* 1991;6:863-866
 84. Shirsat NV, Pignolo RJ, Kreider BL, Rovera G. A member of the ras gene superfamily is expressed specifically in T, B and myeloid hemopoietic cells *Oncogene* 1990;5:769-772
 85. Haataja L, Groffen J, Heisterkamp N. Characterization of RAC3, a novel member of the Rho family *J. Biol. Chem.* 1997;272:20384-20388
 86. Gu Y, Filippi MD, Cancelas JA et al. Hematopoietic cell regulation by Rac1 and Rac2 guanosine triphosphatases *Science* 2003;302:445-449
 87. Gao Y, Dickerson JB, Guo F, Zheng J, Zheng Y. Rational design and characterization of a Rac GTPase-specific small molecule inhibitor *Proc. Natl. Acad. Sci. U.S.A* 2004;101:7618-7623
 88. Muller LU, Schore RJ, Zheng Y, Thomas EK, Kim MO, Cancelas JA, Gu Y, Williams DA. Rac guanosine triphosphatases represent a potential target in AML *Leukemia* 2008;22:1803-1806
 89. Rozenveld-Geugien M, Baas IO, van Gosliga D, Vellenga E, Schuringa JJ. Expansion of normal and leukemic human hematopoietic stem/progenitor cells requires rac-mediated interaction with stromal cells *Exp. Hematol.* 2007;35:782-792
 90. Beachy PA, Karhadkar SS, Berman DM. Tissue repair and stem cell renewal in carcinogenesis *Nature* 2004;432:324-331
 91. Rohatgi R and Scott MP. Patching the gaps in Hedgehog signalling *Nat. Cell Biol.* 2007;9:1005-1009
-

92. Duman-Scheel M, Weng L, Xin S, Du W. Hedgehog regulates cell growth and proliferation by inducing Cyclin D and Cyclin E *Nature* 2002;417:299-304
93. Liu S, Dontu G, Mantle ID, Patel S, Ahn NS, Jackson KW, Suri P, Wicha MS. Hedgehog signaling and Bmi-1 regulate self-renewal of normal and malignant human mammary stem cells *Cancer Res.* 2006;66:6063-6071
94. Chen JK, Taipale J, Cooper MK, Beachy PA. Inhibition of Hedgehog signaling by direct binding of cyclopamine to Smoothened *Genes Dev.* 2002;16:2743-2748
95. Zhao C, Chen A, Jamieson CH, Fereshteh M, Abrahamsson A, Blum J, Kwon HY, Kim J, Chute JP, Rizzieri D, Munchhof M, Vanarsdale T, Beachy PA, Reya T. Hedgehog signalling is essential for maintenance of cancer stem cells in myeloid leukaemia *Nature* 2009;458:776-779
96. Reya T, Duncan AW, Ailles L, Domen J, Scherer DC, Willert K, Hintz L, Nusse R, Weissman IL. A role for Wnt signalling in self-renewal of haematopoietic stem cells *Nature* 2003;423:409-414
97. Reya T and Clevers H. Wnt signalling in stem cells and cancer *Nature* 2005;434:843-850
98. Giles RH, van Es JH, Clevers H. Caught up in a Wnt storm: Wnt signaling in cancer *Biochim.Biophys.Acta* 2003;1653:1-24
99. He TC, Sparks AB, Rago C, Hermeking H, Zawel L, da Costa LT, Morin PJ, Vogelstein B, Kinzler KW. Identification of c-MYC as a target of the APC pathway *Science* 1998;281:1509-1512
100. Tetsu O and McCormick F. Beta-catenin regulates expression of cyclin D1 in colon carcinoma cells *Nature* 1999;398:422-426
101. Wielenga VJ, Smits R, Korinek V, Smit L, Kielman M, Fodde R, Clevers H, Pals ST. Expression of CD44 in Apc and Tcf mutant mice implies regulation by the WNT pathway *Am.J.Pathol.* 1999;154:515-523
102. Muller-Tidow C, Steffen B, Cauvet T, Tickenbrock L, Ji P, Diederichs S, Sargin B, Kohler G, Stelljes M, Puccetti E, Ruthardt M, deVos S, Hiebert SW, Koeffler HP, Berdel WE, Serve H. Translocation products in acute myeloid leukemia activate the Wnt signaling pathway in hematopoietic cells *Mol.Cell Biol.* 2004;24:2890-2904
103. Mikesch JH, Steffen B, Berdel WE, Serve H, Muller-Tidow C. The emerging role of Wnt signaling in the pathogenesis of acute myeloid leukemia *Leukemia* 2007;21:1638-1647
104. Zhao C, Blum J, Chen A, Kwon HY, Jung SH, Cook JM, Lagoo A, Reya T. Loss of beta-catenin impairs the renewal of normal and CML stem cells in vivo *Cancer Cell* 2007;12:528-541
105. Hu Y, Chen Y, Douglas L, Li S. beta-Catenin is essential for survival of leukemic stem cells insensitive to kinase inhibition in mice with BCR-ABL-induced chronic myeloid leukemia *Leukemia* 2009;23:109-116
106. Kawano Y and Kypta R. Secreted antagonists of the Wnt signalling pathway *J.Cell Sci.* 2003;116:2627-2634
107. Zhu Y, Sun Z, Han Q, Liao L, Wang J, Bian C, Li J, Yan X, Liu Y, Shao C, Zhao RC. Human mesenchymal stem cells inhibit cancer cell proliferation by secreting DKK-1 *Leukemia* 2009;23:925-933
108. Lepourcelet M, Chen YN, France DS, Wang H, Crews P, Petersen F, Bruseo C, Wood AW, Shivdasani RA. Small-molecule antagonists of the oncogenic Tcf/beta-catenin protein complex *Cancer Cell* 2004;5:91-102
109. Minke KS, Staib P, Puetter A et al. Small molecule inhibitors of WNT signaling effectively induce apoptosis in acute myeloid leukemia cells *Eur.J.Haematol.* 2009;82:165-175
110. Sophos NA and Vasilou V. Aldehyde dehydrogenase gene superfamily: the 2002 update *Chem.Biol.Interact.* 2003;143-144:5-22
111. Moreb JS, Mohuczy D, Ostmark B, Zucali JR. RNAi-mediated knockdown of aldehyde dehydrogenase class-1A1 and class-3A1 is specific and reveals that each contributes equally to the resistance against 4-hydroperoxycyclophosphamide *Cancer Chemother.Pharmacol.* 2007;59:127-136
112. Pearce DJ, Taussig D, Simpson C, Allen K, Rohatiner AZ, Lister TA, Bonnet D. Characterization of cells with a high aldehyde dehydrogenase activity from cord blood and acute myeloid leukemia samples *Stem Cells* 2005;23:752-760
113. Ginestier C, Hur MH, Charafe-Jauffret E, Monville F, Dutcher J, Brown M, Jacquemier J, Viens P, Kleer CG, Liu S, Schott A, Hayes D, Birnbaum D, Wicha MS, Dontu G. ALDH1 is a marker of normal and malignant human mammary stem cells and a predictor of poor clinical outcome *Cell Stem Cell* 2007;1:555-567

114. Ma S, Chan KW, Lee TK, Tang KH, Wo JY, Zheng BJ, Guan XY. Aldehyde dehydrogenase discriminates the CD133 liver cancer stem cell populations *Mol.Cancer Res.* 2008;6:1146-1153
115. Cheung AM, Wan TS, Leung JC, et al. Aldehyde dehydrogenase activity in leukemic blasts defines a subgroup of acute myeloid leukemia with adverse prognosis and superior NOD/SCID engrafting potential *Leukemia* 2007;21:1423-1430
116. Dylla SJ, Beviglia L, Park IK et al. Colorectal cancer stem cells are enriched in xenogeneic tumors following chemotherapy *PLoS.ONE.* 2008;3:e2428-
117. Rynningen A, Stapnes C, Paulsen K, Lassalle P, Gjertsen BT, Bruserud O. In vivo biological effects of ATRA in the treatment of AML *Expert.Opin.Investig.Drugs* 2008;17:1623-1633
118. Dahia PL, Aguiar RC, Alberta J et al. PTEN is inversely correlated with the cell survival factor Akt/PKB and is inactivated via multiple mechanisms in haematological malignancies *Hum.Mol.Genet.* 1999;8:185-193
119. Cheong JW et al. Phosphatase and tensin homologue phosphorylation in the C-terminal regulatory domain is frequently observed in acute myeloid leukaemia and associated with poor clinical outcome *Br.J.Haematol.* 2003;122:454-456
120. Maehama T and Dixon JE. The tumor suppressor, PTEN/MMAC1, dephosphorylates the lipid second messenger, phosphatidylinositol 3,4,5-trisphosphate *J.Biol.Chem.* 1998;273:13375-13378
121. Yilmaz OH, Valdez R, Theisen BK, Guo W, Ferguson DO, Wu H, Morrison SJ. Pten dependence distinguishes haematopoietic stem cells from leukaemia-initiating cells *Nature* 2006;441:475-482
122. Ito K, Bernardi R, Morotti A, Matsuoka S, Saglio G, Ikeda Y, Rosenblatt J, Avigan DE, Teruya-Feldstein J, Pandolfi PP. PML targeting eradicates quiescent leukaemia-initiating cells *Nature* 2008;453:1072-1078
123. Martinez A, et al. First non-ATP competitive glycogen synthase kinase 3 beta (GSK-3beta) inhibitors: thiadiazolidinones (TDZD) as potential drugs for the treatment of Alzheimer's disease *J.Med.Chem.* 2002;45:1292-1299
124. Abrahamsson AE, Geron I, Gotlib J, Dao KH, Barroga CF, Newton IG, Giles FJ, Durocher J, Creusot RS, Karimi M, Jones C, Zehnder JL, Keating A, Negrin RS, Weissman IL, Jamieson CH. Glycogen synthase kinase 3beta missplicing contributes to leukemia stem cell generation *Proc.Natl.Acad.Sci.U.S.A* 2009;106:3925-3929
125. Guzman ML et al.. Rapid and selective death of leukemia stem and progenitor cells induced by the compound 4-benzyl, 2-methyl, 1,2,4-thiadiazolidine, 3,5 dione (TDZD-8) *Blood* 2007;110:4436-4444
126. Steele AJ, Jones DT, Ganeshaguru K, Duke VM, Yogashangary BC, North JM, Lowdell MW, Kottaridis PD, Mehta AB, Prentice AG, Hoffbrand AV, Wickremasinghe RG. The sesquiterpene lactone parthenolide induces selective apoptosis of B-chronic lymphocytic leukemia cells in vitro *Leukemia* 2006;20:1073-1079
127. Wen J, You KR, Lee SY, Song CH, Kim DG. Oxidative stress-mediated apoptosis. The anticancer effect of the sesquiterpene lactone parthenolide *J.Biol.Chem.* 2002;277:38954-38964
128. Oka D, Nishimura K, Shiba M, Nakai Y, Arai Y, Nakayama M, Takayama H, Inoue H, Okuyama A, Nonomura N. Sesquiterpene lactone parthenolide suppresses tumor growth in a xenograft model of renal cell carcinoma by inhibiting the activation of NF-kappaB *Int.J.Cancer* 2007;120:2576-2581
129. Guzman ML, Rossi RM, Karnischky L, Li X, Peterson DR, Howard DS, Jordan CT. The sesquiterpene lactone parthenolide induces apoptosis of human acute myelogenous leukemia stem and progenitor cells *Blood* 2005;105:4163-4169
130. Guzman ML, Rossi RM, Neelakantan S, Li X, Corbett CA, Hassane DC, Becker MW, Bennett JM, Sullivan E, Lachowicz JL, Vaughan A, Sweeney CJ, Matthews W, Carroll M, Liesveld JL, Crooks PA, Jordan CT. An orally bioavailable parthenolide analog selectively eradicates acute myelogenous leukemia stem and progenitor cells *Blood* 2007;110:4427-4435
131. Hanahan D and Weinberg RA. The hallmarks of cancer *Cell* 2000;100:57-70
132. Jones PA and Baylin SB. The epigenomics of cancer *Cell* 2007;128:683-692
133. Ringrose L and Paro R. Epigenetic regulation of cellular memory by the Polycomb and Trithorax group proteins *Annu. Rev.Genet.* 2004;38:413-443
134. Schuebel K et al. CIMP origin for promoter hypermethylation in colorectal cancer? *Nat.Genet.* 2006;38:738-740
135. Widschwendter M, Fiegler H, Egle D, Mueller-Holzner E, Spizzo G, Marth C, Weisenberger DJ, Campan M, Young J, Jacobs I, Laird PW. Epigenetic stem cell signature in cancer *Nat.Genet.* 2007;39:157-158

136. Park IK, Qian D, Kiel M, Becker MW, Pihalja M, Weissman IL, Morrison SJ, Clarke MF. Bmi-1 is required for maintenance of adult self-renewing haematopoietic stem cells *Nature* 2003;423:302-305
137. Chowdhury M, Mihara K, Yasunaga S, Ohtaki M, Takiyama Y, Kimura A. Expression of Polycomb-group (PcG) protein BMI-1 predicts prognosis in patients with acute myeloid leukemia *Leukemia* 2007;21:1116-1122
138. Mohty M, Yong AS, Szydlowski RM, Apperley JF, Melo JV. The polycomb group BMI1 gene is a molecular marker for predicting prognosis of chronic myeloid leukemia *Blood* 2007;110:380-383
139. Wang GG, Song J, Wang Z, Dormann HL, Casadio F, Li H, Luo JL, Patel DJ, Allis CD. Haematopoietic malignancies caused by dysregulation of a chromatin-binding PHD finger *Nature* 2009;459:847-851
140. Taverna SD, Li H, Ruthenburg AJ, Allis CD, Patel DJ. How chromatin-binding modules interpret histone modifications: lessons from professional pocket pickers *Nat.Struct.Mol.Biol.* 2007;14:1025-1040
141. Baker LA, Allis CD, Wang GG. PHD fingers in human diseases: disorders arising from misinterpreting epigenetic marks *Mutat.Res.* 2008;647:3-12
142. van Zutven LJ, Onen E, Velthuisen SC, van Drunen E, von Bergh AR, van den Heuvel-Eibrink MM, Veronese A, Mecucci C, Negrini M, de Greef GE, Beverloo HB. Identification of NUP98 abnormalities in acute leukemia: JARID1A (12p13) as a new partner gene *Genes Chromosomes.Cancer* 2006;45:437-446
143. Wang GG, Cai L, Pasillas MP, Kamps MP. NUP98-NSD1 links H3K36 methylation to Hox-A gene activation and leukaemogenesis *Nat.Cell Biol.* 2007;9:804-812
144. Moore MA, Chung KY, Plasilova M, Schuringa JJ, Shieh JH, Zhou P, Morrone G. NUP98 dysregulation in myeloid leukemogenesis *Ann.N.Y.Acad.Sci.* 2007;1106:114-142
145. Wang GG, Song J, Wang Z, Dormann HL, Casadio F, Li H, Luo JL, Patel DJ, Allis CD. Haematopoietic malignancies caused by dysregulation of a chromatin-binding PHD finger *Nature* 2009;
146. Lessard J and Sauvageau G. Bmi-1 determines the proliferative capacity of normal and leukaemic stem cells *Nature* 2003;423:255-260
147. Liang G, Gonzales FA, Jones PA, Orntoft TF, Thykjaer T. Analysis of gene induction in human fibroblasts and bladder cancer cells exposed to the methylation inhibitor 5-aza-2'-deoxycytidine *Cancer Res.* 2002;62:961-966
148. Kuendgen A and Gattermann N. Valproic acid for the treatment of myeloid malignancies *Cancer* 2007;110:943-954
149. Guo F, Sigua C, Tao J, Bali P, George P, Li Y, Wittmann S, Moscinski L, Atadja P, Bhalla K. Cotreatment with histone deacetylase inhibitor LAQ824 enhances Apo-2L/tumor necrosis factor-related apoptosis inducing ligand-induced death inducing signaling complex activity and apoptosis of human acute leukemia cells *Cancer Res.* 2004;64:2580-2589
150. Morotti A, Cilloni D, Mesa F, Arruga F, Defilippi I, Carturan S, Catalano R, Rosso V, Chiarenza A, Pilatino C, Guerrasio A, Tauli R, Bracco E, Pautasso M, Baraban D, Gottardi E, Saglio G. Valproate enhances imatinib-induced growth arrest and apoptosis in chronic myeloid leukemia cells *Cancer* 2006;106:1188-1196
151. Yu C, Rahmani M, Conrad D et al. The proteasome inhibitor bortezomib interacts synergistically with histone deacetylase inhibitors to induce apoptosis in Bcr/Abl+ cells sensitive and resistant to STI571 *Blood* 2003;102:3765-3774
152. Saito Y, Liang G, Egger G, Friedman JM, Chuang JC, Coetzee GA, Jones PA. Specific activation of microRNA-127 with downregulation of the proto-oncogene BCL6 by chromatin-modifying drugs in human cancer cells *Cancer Cell* 2006;9:435-443
